

1-24: Calculate y' :

1. $y = (x^2 + x^3)^4$

$y' = 4(x^2 + x^3)^3 [2x + 3x^2]$

$y' = 4(2x + 3x^2)(x^2 + x^3)^3$

2. $y = \frac{1}{\sqrt{x}} - \frac{1}{\sqrt[5]{x^3}}$

$y = x^{-1/2} - x^{-3/5}$

$y = -\frac{1}{2}x^{-3/2} + \frac{3}{5}x^{-8/5}$

$y' = -\frac{1}{2x^{3/2}} + \frac{3}{5x^{8/5}}$

3. $y = \frac{x^2 - x - 2}{\sqrt{x}}$

$y = \frac{x^2}{x^{1/2}} - \frac{x}{x^{1/2}} - \frac{2}{x^{1/2}}$

$y = x^{3/2} - x^{1/2} - 2x^{-1/2}$

$y' = \frac{3}{2}x^{1/2} - \frac{1}{2}x^{-1/2} + x^{-3/2}$

$y' = \frac{3x^{1/2}}{2} - \frac{1}{2x^{1/2}} + \frac{1}{x^{3/2}}$

4. $y = \frac{\tan x}{1 + \cos x}$

$y' = \frac{(1 + \cos x)\sec^2 x - \tan x(\sin x)}{(1 + \cos x)^2}$

$y' = \frac{\sec^2 x(1 + \cos x) + \sin x \tan x}{(1 + \cos x)^2}$

Can keep going \therefore look at last page if interested

5. $y = x^2 \sin(\pi x)$

$y' = x^2 [\cos(\pi x) \cancel{\pi}] + \sin(\pi x) [2x]$

$y' = \pi x^2 \cos(\pi x) + 2x \sin(\pi x)$

6. $y = \frac{t^4 - 1}{t^4 + 1}$

$y' = \frac{(t^4 + 1)(4t^3) - (t^4 - 1)(4t^3)}{(t^4 + 1)^2}$

$y' = 4t^1 + 4t^3 - 4t^1 + 4t^3$

$y' = \frac{8t^3}{(t^4 + 1)^2}$

7. $y = \ln(x \ln x)$

$\frac{1}{x \ln x} \frac{d}{dx} [x \ln x]$

$\frac{1}{x \ln x} \left[x \left(\frac{1}{x} \right) + \ln x (1) \right]$

$\frac{1}{x \ln x} [1 + \ln x]$

$\frac{1 + \ln x}{x \ln x}$

8. $y = \sqrt{x} \cos \sqrt{x} = x^{1/2} \cos x^{1/2}$

$y' = x^{1/2} (-\sin x^{1/2}) \frac{d}{dx} [x^{1/2}]$

$+ \cos x^{1/2} \left[\frac{1}{2}x^{-1/2} \right]$

$y' = -\sqrt{x} \sin \sqrt{x} \left(\frac{1}{2}x^{-1/2} \right)$

$+ \cos \sqrt{x} \left(\frac{1}{2\sqrt{x}} \right)$

$y' = -\frac{\sqrt{x} \sin \sqrt{x}}{2\sqrt{x}} + \frac{\cos \sqrt{x}}{2\sqrt{x}}$

$= -\frac{\sqrt{x} \sin \sqrt{x} + \cos \sqrt{x}}{2\sqrt{x}}$

9. $y = \frac{e^x}{x^2}$

$x^2 e^{\frac{1}{x}} \cdot \frac{d}{dx} [x^{-2}] - e^{\frac{1}{x}} (2x)$

$x^2 e^{\frac{1}{x}} (-x^{-2} x^4) - 2x e^{\frac{1}{x}}$

$x^2 e^{\frac{1}{x}} \left(-\frac{x^4}{x^2}\right) - 2x e^{\frac{1}{x}}$

$-e^{\frac{1}{x}} - 2x e^{\frac{1}{x}} = \frac{-e^{\frac{1}{x}} (1+2x)}{x^4}$

$$10. y = \ln \sec x$$

$$\frac{1}{\sec x} \frac{d}{dx} [\sec x]$$

$$\frac{1}{\sec x} [\sec x \tan x]$$

$$= \tan x$$

$$11. y = \cot(\csc x)$$

$$-\csc^2(\csc x) \frac{d}{dx} [\csc x]$$

$$-\csc^2(\csc x) [-\csc x \cot x]$$

$$\csc^2(\csc x) \csc x \cot x$$

$$12. y = e^{x \sec x}$$

$$e^{x \sec x} \frac{d}{dx} [x \sec x]$$

$$e^{x \sec x} [x \sec x \tan x + \sec x (1)]$$

$$e^{x \sec x} [x \sec x \tan x + \sec x]$$

$$e^{x \sec x} \cdot \sec x [x \tan x + 1]$$

$$13. y = 3^{x \ln x}$$

$$y' = 3^{x \ln x} \cdot \ln 3 \cdot \frac{d}{dx} [x \ln x]$$

$$= 3^{x \ln x} \cdot \ln 3 \left[x \cdot \frac{1}{x} + \ln x (1) \right]$$

$$= 3^{x \ln x} \cdot \ln 3 [1 + \ln x]$$

$$14. y = \sec(1+x^2)$$

$$\sec(1+x^2) \tan(1+x^2) [2x]$$

$$2x \sec(1+x^2) \tan(1+x^2)$$

$$15. y = \log_5(1+2x)$$

$$y' = \frac{1}{(1+2x) \ln 5} [2]$$

$$= \frac{2}{(1+2x) \ln 5}$$

$$16. y = (\cos x)^x$$

$$\ln y = x \cdot \ln \cos x$$

$$\frac{1}{y} [y'] = x \left(\frac{1}{\cos x} (-\sin x) + \ln \cos x (1) \right)$$

$$\frac{1}{y} [y'] = [-x \tan x + \ln \cos x] \textcolor{red}{y}$$

$$y' = [-x \tan x + \ln \cos x] \cos x^x$$

$$17. y = \frac{(x^2+1)^4}{(2x+1)^3 (3x-1)^5}$$

$$\ln y = 4 \ln(x^2+1) - 3 \ln(2x+1) - 5 \ln(3x-1)$$

$$\frac{1}{y} [y'] = 4 \left(\frac{1}{x^2+1} \right) (2x) - 3 \left(\frac{1}{2x+1} \right) (2) - 5 \left(\frac{1}{3x-1} \right) (3)$$

$$\frac{1}{y} [y'] = \frac{8x}{x^2+1} - \frac{6}{2x+1} - \frac{15}{3x-1}$$

$$y' = \left[\frac{8x}{x^2+1} - \frac{6}{2x+1} - \frac{15}{3x-1} \right] \left[\frac{(x^2+1)^4}{(2x+1)^3 (3x-1)^5} \right]$$

$$18. y = e^{\cos x} + \cos(e^x)$$

$$e^{\cos x} (-\sin x) + (-\sin e^x) e^x$$

$$-\sin x e^{\cos x} - \sin e^x \cdot e^x$$

$$19. y = 10^{\tan \pi \theta}$$

$$\begin{aligned}y' &= 10^{\tan \pi \theta} \cdot \ln 10 \cdot \frac{d}{d\theta}(\tan \pi \theta) - \csc^2(3x^2+5)[6x] \\&= 10^{\tan \pi \theta} \cdot \ln 10 \cdot \sec^2 \pi \theta \frac{d}{d\theta}(\pi \theta) - 6x \csc^2(3x^2+5) \\&= \pi 10^{\tan \pi \theta} \cdot \ln 10 \cdot \sec^2 \pi \theta\end{aligned}$$

$$20. y = \cot(3x^2 + 5)$$

$$- \csc^2(3x^2+5)[6x]$$

$$\begin{aligned}21. y &= \sqrt{t \ln(t^4)} = (t \ln(t^4))^{1/2} \\&\frac{1}{2} [t \ln(t^4)]^{-1/2} \left[t \left(\frac{1}{t^4} \right) (4t^3) + \ln(t^4)(1) \right] \\&\frac{1}{2} [t \ln(t^4)]^{-1/2} [4 + \ln(t^4)] \\&\frac{4 + \ln(t^4)}{2 \sqrt{t \ln(t^4)}}\end{aligned}$$

$$22. y = \tan^2(\sin \theta) = [\tan(\sin \theta)]^2$$

$$2[\tan(\sin \theta)] \frac{d}{d\theta} [\tan(\sin \theta)]$$

$$2\tan(\sin \theta) \sec^2(\sin \theta) \frac{d}{d\theta} [\sin \theta]$$

$$2\tan(\sin \theta) \sec^2(\sin \theta) \cos \theta$$

$$23. y = (3x+2)^5(5-x)^8$$

$$(3x+2)^5(8)(5-x)^7(-1) + (5-x)^8(3x+2)^4(-8(3x+2)^5(5-x)^7) + 15(5-x)^8(3x+2)^4$$

$$(3x+2)^4(5-x)^7[-8(3x+2)+15(5-x)]$$

$$(3x+2)^4(5-x)^7[-24x-16+75-15x]$$

$$(3x+2)^4(5-x)^7[-39x+59]$$

$$24. y = \left(\frac{2x-5}{x+4} \right)^8$$

$$8 \left(\frac{2x-5}{x+4} \right)^7 \left[\frac{(x+4)(2)-(2x-5)(1)}{(x+4)^2} \right]$$

$$8(2x-5)^7 [2x+8-2x+5] \\ (x+4)^7 (x+4)^2$$

$$\frac{8(2x-5)^7(13)}{(x+4)^9} = \frac{104(2x-5)^7}{(x+4)^9}$$

25-27: Find an equation of the tangent to the curve at the given point.

$$25. y = 4 \sin^2 x, \quad \left(\frac{\pi}{6}, 1 \right)$$

$$y' = 8 \sin x \cos x$$

$$y'\left(\frac{\pi}{6}\right) = 8\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) = 2\sqrt{3}$$

$$y-1 = 2\sqrt{3}\left(x - \frac{\pi}{6}\right)$$

$$26. y = \frac{x^2-1}{x^2+1}, \quad (0, -1)$$

$$y' = \frac{(x^2+1)(2x) - (x^2-1)(2x)}{(x^2+1)^2}$$

$$y' = \frac{2x^3 + 2x - 2x^3 + 2x}{(x^2+1)^2} = \frac{4x}{(x^2+1)^2}$$

$$y'(0) = \frac{4(0)}{(0^2+1)^2} = 0$$

$$y-(-1) = 0(x-0) \quad [y=-1]$$

$$27. y = \sqrt{1+4 \sin x}, \quad (0, 1)$$

$$y = (1+4 \sin x)^{1/2}$$

$$y' = \frac{1}{2}(1+4 \sin x)^{-1/2} \cdot 4 \cos x$$

$$y' = \frac{4 \cos x}{2\sqrt{1+4 \sin x}}$$

$$y'(0) = \frac{4 \cos(0)}{2\sqrt{1+4 \sin(0)}} = \frac{4(1)}{2\sqrt{1}} = 2$$

28-29: Suppose that $h(x) = f(x)g(x)$ and $F(x) = f(g(x))$, where $f(2) = 3$, $g(2) = 5$, $g'(2) = 4$, $f'(2) = -2$, and $f'(5) = 11$.

28. Find $h'(2)$

$$h(x) = f(x)g(x)$$

$$h'(x) = f(x)g'(x) + g(x)f'(x)$$

$$h'(2) = f(2)g'(2) + g(2)f'(2)$$

$$= (3)(4) + (5)(-2)$$

$$= 12 - 10$$

$$= 2$$

29. Find $F'(2)$

$$F(x) = f(g(x))$$


$$F'(x) = f'(g(x)) \cdot g'(x)$$

$$F'(2) = f'[g(2)] \cdot g'(2)$$

$$F'(2) = f'[5][4] \\ (11)(4)$$

30-32: Suppose f and g are functions whose graphs are shown, let $P(x) = f(x)g(x)$, $Q(x) = f(x)/g(x)$, and $C(x) = f(g(x))$.

$$Q(x) = \frac{f(x)}{g(x)}$$



30. Find $P'(2)$

$$\begin{aligned} P(x) &= f(x)g(x) \\ P'(x) &= f(x)g'(x) + g(x)f'(x) \\ P'(2) &= f(2)g'(2) + g(2)f'(2) \\ P'(2) &= (1)(2) + (4)(-1) \\ P'(2) &= 2 - 4 \\ P'(2) &= -2 \end{aligned}$$

31. Find $Q'(2)$

$$\begin{aligned} Q'(x) &= \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2} \\ Q'(2) &= \frac{g(2)f'(2) - f(2)g'(2)}{[g(2)]^2} \\ Q'(2) &= \frac{4(-1) - (1)(2)}{4^2} \\ Q'(2) &= -\frac{6}{16} = -\frac{3}{8} \end{aligned}$$

32. Find $C'(2)$

$$\begin{aligned} C(x) &= f(g(x)) \\ C'(x) &= f'(g(x)) \cdot g'(x) \\ C'(2) &= f'(g(2)) \cdot g'(2) \\ C'(2) &= f'(4) \cdot [2] \\ C'(2) &= [3][2] \\ C'(2) &= 6 \end{aligned}$$

33-40: Find f' in terms of g'

33. $f(x) = x^2g(x)$

$$\begin{aligned} x^2g'(x) + g(x)[2x] \\ x^2g'(x) + 2xg(x) \end{aligned}$$

34. $f(x) = g(x^2)$

$$\begin{aligned} g'(x^2)[2x] \\ 2xg'(x^2) \end{aligned}$$

35. $f(x) = [g(x)]^2$

$$\begin{aligned} 2[g(x)]'g'(x) \\ 2g(x)g'(x) \end{aligned}$$

36. $f(x) = g(g(x))$

$$g'[g(x)] \cdot g'(x)$$

37. $f(x) = g(e^x)$

$$g'[e^x] \cdot e^x$$

38. $f(x) = e^{g(x)}$

$$e^{g(x)} \cdot g'(x)$$

39. $f(x) = \ln|g(x)|$

$$\begin{aligned} \frac{1}{g(x)} \cdot g'(x) \\ \frac{g'(x)}{g(x)} \end{aligned}$$

40. $f(x) = g(\ln x)$

$$\begin{aligned} g'[\ln x] \cdot \frac{1}{x} \\ \frac{g'[\ln x]}{x} \end{aligned}$$